



Contribution ID : 57

Type : Oral

## Overcoming Fermionic Sign Problem in Lattice Quantum Monte Carlo: A Cuprate Case

Wednesday, 25 September 2024 11:55 (25)

We developed a strong coupling perturbation scheme for general Hubbard model around half-field particle-hole symmetric reference system [1]. The approach based on a lattice determinantal Quantum Monte Carlo method in continuous and discrete time versions [2] for a large periodic clusters in a fermionic bath. The first and second order perturbation in the shift of chemical potential and long-range hopping gives a reasonable accuracy for parameters corresponding to the optimal cuprate systems. We calculate spectral function of doped  $t$ - $t'$ - $U$  model for interaction strength equal to the band width and discuss a mechanism of the pseudogap formation. Results for standard cuprates model with  $U=8t=W$  and  $t'/t=-0.3$  for the temperature of the order of  $T=0.1t$  show formation of the Fermi-arcs. We discuss the magnetic and superconducting instability using symmetry broken external fields.

[1] S. Iskakov, M. I. Katsnelson, A. I. Lichtenstein, npj Comp. Materials 10, 36 (2024).

[2] E. Gull, A. J. Millis, A. I. Lichtenstein, et. al., Rev. Mod. Phys. 83, 349 (2011).

Primary author(s) : LICHTENSTEIN, Alexander (University of Hamburg)

Presenter(s) : LICHTENSTEIN, Alexander (University of Hamburg)

Session Classification : Session